A NOTE ON MEASURE EXTENSION PROBLEM OF ℓ -GROUP-VALUED MEASURES

SURJIT SINGH KHURANA

ABSTRACT. For an Abelian, Archimedean Dedekind complete lattice-ordered, weakly σ -distributive group G and an algebra $\mathcal{F}$ of subsets of a set X , every positive measure $\mu: \mathcal{F} \rightarrow G$ has a unique countably additive (in order convergence) extension to σ -algebra generated by $\mathcal{F}$.

1. Introduction and notations

In this note, R is the set of real numbers and $\bar{R} = [-\infty, \infty]$, G is an Abelian, Archimedean Dedekind complete lattice-ordered, weakly σ -distributive group, $\mathcal{F}$ is an algebra of subsets of a set X and $\mu: \mathcal{F} \rightarrow G$ a positive measure (this means for a disjoint sequence $\{A_n\} \subset \mathcal{F}$ with $\cup A_n \in \mathcal{F}$, $\mu(A) = \sum \mu(A_n)$, considering limit in order convergence) and $\mathcal{A}$ the σ -algebra generated by $\mathcal{F}$.

In [6], using Carathéodory measurability, the author has given a very interesting sophisticated proof of the unique extension, in order convergence, of the measure to $\mathcal{A}$ (in [4, 5] the author has given other proofs of this result). In this note we prove that this result can also be easily obtained from the representation theorem proved in [8] (see also [3]).

2. Main result

THEOREM 1 ([6, Theorem 5.1]). *Let $\mu: \mathcal{F} \rightarrow G$ be a positive measure. Then there is a unique extension $\mu: \mathcal{A} \rightarrow G$ which is positive and countably additive in order convergence.*

Proof. For a compact Hausdorff Stonian (extremely disconnected) space Y , $\mathcal{E}(Y)$ will denote the space of all continuous functions $f: Y \rightarrow \bar{R}$ such that f is finite-valued almost everywhere (this means the set where it is $\pm\infty$ is nowhere dense). $\square$

As in [8], we can assume, that $\mathcal{F}$ separates the points of X . As in [8, p. 70], there is a totally disconnected compact Hausdorff space $\tilde{X}$, the Boolean space for $\mathcal{F}$, in which X is dense. If $\tilde{\mathcal{F}}$ and $\mathcal{B}$ are the classes of all clopen subsets and all Baire subsets of $\tilde{X}$, then the mapping $\rho: B \rightarrow B \cap X$, when restricted to $\tilde{\mathcal{F}}$, is an isomorphism onto $\mathcal{F}$ and when restricted to $\mathcal{B}$ is a σ -homomorphism of $\mathcal{B}$ onto a class which contains $\mathcal{A}$ with kernel being the class of Baire sets disjoint from X [8]. The measure μ gives a finitely additive positive mapping

$$\tilde{\mu}: \tilde{\mathcal{F}} \rightarrow G, \quad \tilde{\mu}(A) = \mu(A \cap X).$$

By [2, Theorem 4], there is a compact Hausdorff Stonian space Y such that G is lattice group isomorphic to a sub-lattice group of $\mathcal{E}(Y)$, preserving arbitrary sup and inf; we denote this isomorphism by i . Put $i \circ \mu(X) = e_0$ and let Y_0 be an open dense subset of Y where e_0 is finite-valued. We denote by $C(Y_0)$ the space of all real-valued continuous functions on the locally compact space Y_0 . The mapping $C(Y_0) \rightarrow \mathcal{E}(Y)$, $f \rightarrow \hat{f}$ (continuous extension) is a lattice isomorphism preserving arbitrary sup and inf. Put $e = (e_0)_{|Y_0}$. The finitely additive positive measure

$$\nu: \tilde{\mathcal{F}} \rightarrow C(Y_0)$$

is defined as

$$\nu(A) = (i \circ \tilde{\mu}(A))_{|Y_0}.$$

Putting

$$I = \{f \in C(Y_0) : |f| \leq e\},$$

we get a Dedekind complete vector sub-lattice E of $C(Y_0)$, $E = \bigcup_{n=1}^{\infty} nI$ with a strong unit e and a positive finitely additive measure $\nu: \tilde{\mathcal{F}} \rightarrow E$. Defining the norm $\|\cdot\|$ on E , $\|f\| = \inf\{\lambda > 0 : |f| \leq \lambda e\}$, E becomes a Dedekind complete M -space [1] and so there is an extremely disconnected compact Hausdorff space S such that E is linear lattice isomorphic to $C(S)$, preserving arbitrary sup and inf. Thus we get a finitely additive positive measure $\nu: \tilde{\mathcal{F}} \rightarrow C(S)$ which extends to linear positive mapping

$$\nu: \mathcal{S}(\tilde{\mathcal{F}}) \rightarrow C(S), \quad \mathcal{S}(\tilde{\mathcal{F}})$$

being the space of all real-valued $\tilde{\mathcal{F}}$ -simple functions on X ; with sup norm topology on $\mathcal{S}(\tilde{\mathcal{F}})$, this mapping is easily verified to be continuous. Since X is totally disconnected, by Stone-Weierstrass theorem, $\mathcal{S}(\tilde{\mathcal{F}})$ is norm dense in $C(\tilde{X})$. Thus, we get a positive continuous mapping

$$\nu: C(\tilde{X}) \rightarrow C(S).$$

MEASURE EXTENSION

By [7], we get a quasi-regular Baire measure on $\tilde{X}$ which extends ν . Since G is weakly σ -distributive, the measure $\nu: \mathcal{B} \rightarrow G$ is regular. Now, for any compact G_δ -set $C \subset \tilde{X} \setminus X$, there is a decreasing sequence $\{F_n\} \subset \tilde{\mathcal{F}}$ such that $F_n \downarrow \chi_C$; since μ is a measure, we get $\nu(C) = 0$. By regularity, $\nu(B) = 0$ for any Baire subset of $\tilde{X}$ disjoint from X . Now, for any $A \in \mathcal{A}$ we can define $\mu(A) = \nu(B)$ for any Baire subset B of $\tilde{X}$ with $B \cap X = A$. The rest is easily verified.

REFERENCES

- [1] ALIPRANTIS, C. D.—BURKINSHAW, O.: *Positive Operators*. Academic Press, Orlando, 1985.
- [2] BERNAU, S. J.: *Unique representation of Archimedean lattice groups and normal Archimedean lattice rings*, Proc. London Math. Soc. (3) **15** (1965), 599–631.
- [3] KHURANA, S. S.: *Lattice-valued Borel measures*, Rocky Mountain J. Math. **6** (1976), 377–382.
- [4] RIEČAN, B.: *On measures and integrals with values in ordered groups*, Math. Slovaca **33** (1983), 153–163.
- [5] RIEČAN, B.—NEUBRUNN, T.: *Integral, Measure and Ordering*. Math. Appl., Vol. 411, Kluwer Acad. Publ., Dordrecht, 1997.
- [6] RIEČAN, B.: *Carathéodory measurability revisited*, Tatra Mt. Math. Publ. **34** (2006), 321–332.
- [7] WRIGHT J. D. M.: *Stone-algebra-valued measures and integrals*, Proc. London Math. Soc. **19** (1969), 107–122.
- [8] WRIGHT J. D. M.: *The measure problem for vector lattices*, Ann. Inst. Fourier (Grenoble) **21** (1971), 65–85.

Received July 20, 2008

*Department of Mathematics
The University of Iowa
Iowa City, Iowa 52242
U.S.A.
E-mail: khurana@math.uiowa.edu*