PRODUCTS OF INTERNALLY QUASI-CONTINUOUS FUNCTIONS

PAULINA SZCZUKA — MARIOLA MARCINIAK

ABSTRACT. In this paper we characterize the product of internally quasi-continuous functions and we construct a bounded internally quasi-continuous strong Świątkowski function which cannot be written as a finite product of internally strong Świątkowski functions.

1. Preliminaries

The letters $\mathbb{R}$ and $\mathbb{N}$ denote the real line and the set of positive integers, respectively. The symbol $I(a, b)$ denotes an open interval with the endpoints a and b . For each $A \subset \mathbb{R}$ we use the symbols $\text{int } A$, $\text{cl } A$, $\text{bd } A$, and $\text{card } A$ to denote the interior, the closure, the boundary, and the cardinality of A , respectively. We say that a set $A \subset \mathbb{R}$ is *simply open* [1], if it can be written as the union of an open set and a nowhere dense set.

The word function denotes a mapping from $\mathbb{R}$ into $\mathbb{R}$ unless otherwise explicitly stated. The symbol $\mathcal{C}(f)$ stands for the set of all points of continuity of f . We say that f is a *Darboux function*, if it maps the connected sets onto connected sets. We say that f is *cliquish* [11] ($f \in \mathcal{C}_q$), if the set $\mathcal{C}(f)$ is dense in $\mathbb{R}$. We say that f is *internally cliquish* ($f \in \mathcal{C}_{qi}$), if the set $\text{int } \mathcal{C}(f)$ is dense in $\mathbb{R}$. We say that f is *quasi-continuous* in the sense of Kempisty [5] ($f \in \mathcal{Q}$), if for all $x \in \mathbb{R}$ and open sets $U \ni x$ and $V \ni f(x)$, the set $\text{int}(U \cap f^{-1}(V)) \neq \emptyset$. We say that f is *internally quasi-continuous* [8] ($f \in \mathcal{Q}_i$), if it is quasi-continuous and its set of points of discontinuity is nowhere dense; equivalently, f is internally quasi-continuous if $f|_{\text{int } \mathcal{C}(f)}$ is dense in f . We say that x_0 is a point of internal

© 2013 Mathematical Institute, Slovak Academy of Sciences.

2010 Mathematics Subject Classification: Primary 26A21, 54C30; Secondary 26A15, 54C08.

Keywords: quasi-continuous function, internally quasi-continuous function, strong Świątkowski function, internally strong Świątkowski function, product of functions.

Supported by Kazimierz Wielki University.

quasi-continuity of f if and only if there is a sequence $(x_n) \subset \text{int } \mathcal{C}(f)$ such that $x_n \rightarrow x_0$ and $f(x_n) \rightarrow f(x_0)$ (see [8]). We say that f is a *strong Świątkowski function* [6] ($f \in \acute{\mathcal{S}}_s$), if whenever $a, b \in \mathbb{R}$, $a < b$, and $y \in I(f(a), f(b))$, there is $x_0 \in (a, b) \cap \mathcal{C}(f)$ such that $f(x_0) = y$. We say that f is an *internally strong Świątkowski function* [8] ($f \in \acute{\mathcal{S}}_{si}$), if whenever $a, b \in \mathbb{R}$, $a < b$, and $y \in I(f(a), f(b))$, there is

$$x_0 \in (a, b) \cap \text{int } \mathcal{C}(f) \quad \text{such that} \quad f(x_0) = y.$$

Clearly, each strong Świątkowski function has the Darboux property. Moreover, we can easily see that the following inclusions

$$\acute{\mathcal{S}}_{si} \subset \acute{\mathcal{S}}_s \subset \mathcal{Q} \subset \mathcal{C}_q \quad \text{and} \quad \acute{\mathcal{S}}_{si} \subset \mathcal{Q}_i \subset \mathcal{C}_{qi} \subset \mathcal{C}_q$$

are satisfied.

Finally, the symbol $[f = a]$ stands for the set $\{x \in \mathbb{R} : f(x) = a\}$.

2. Introduction

In 1960 S. Marcus remarked that not every function is the product of Darboux functions [9]. The problem of characterizing the class of products of Darboux functions was solved by J. G. Ceder [2], [3]. In 1985 Z. Grande constructed a nonnegative Baire one function which cannot be the product of a finite number of quasi-continuous functions, and asked for characterization of such products [4]. The following theorem (see [7, Theorem III.2.1]) gives an answer to this question.

THEOREM 2.1. *For each function f the following conditions are equivalent:*

- i) *there are quasi-continuous functions g_1 and g_2 with $f = g_1 g_2$,*
- ii) *f is a finite product of quasi-continuous functions,*
- iii) *f is cliquish and the set $[f = 0]$ is simply open.*

In 1996 A. Maliszewski characterized the product of Darboux quasi-continuous functions [7, Theorem III.3.1] and proved that there exists a bounded Darboux quasi-continuous function which cannot be written as the finite product of strong Świątkowski functions [7, Proposition III.4.1]. Ten years later P. Szczuka characterized the product of four and more strong Świątkowski functions [10, Theorem 4.2].

In this paper we characterize the product of internally quasi-continuous functions (Theorem 4.2) and we construct a bounded internally quasi-continuous strong Świątkowski function which cannot be written as the finite product of internally strong Świątkowski functions (Proposition 4.3).

3. Auxiliary lemmas

Lemma 3.1 is due to A. Maliszewski [7, Lemma III.1.10].

LEMMA 3.1. *Let $I = (a, b)$, $\Gamma > 0$ be an extended real number, and $k > 1$. There are functions $g_1, \dots, g_k$ such that $g_1 \dots g_k = 0$ on $\mathbb{R}$ and for $i \in \{1, \dots, k\}$: $\mathbb{R} \setminus \mathcal{C}(g_i) = \text{bd } I$ and $g_i[(a, c)] = g_i[(c, b)] = (-\Gamma, \Gamma)$ for each $c \in I$.*

The proof of Lemma 3.2 can be found in [10, Lemma 3.4].

LEMMA 3.2. *Assume that $F \subset C$ are closed and $\mathcal{J}$ is a family of components of $\mathbb{R} \setminus C$ such that $C \subset \text{cl} \bigcup \mathcal{J}$. There is a family $\mathcal{J}' \subset \mathcal{J}$ such that:*

- i) *for each $J \in \mathcal{J}$, if $F \cap \text{bd } J \neq \emptyset$, then $J \in \mathcal{J}'$,*
- ii) *for each $c \in F$, if c is a right-hand (left-hand) limit point of C , then c is a right-hand (respectively left-hand) limit point of the union $\bigcup \mathcal{J}'$,*
- iii) *$\text{cl} \bigcup \mathcal{J}' \subset F \cup \bigcup_{J \in \mathcal{J}'} \text{cl } J$.*

LEMMA 3.3. *Let $I = (a, b)$ and let the function $f: \text{cl } I \rightarrow (0, +\infty)$ be continuous. There are continuous functions $\psi_1, \psi_2: I \rightarrow (0, +\infty)$ such that $f = \psi_1 \psi_2$ on I and $\psi_i[(a, c)] = \psi_i[(c, b)] = (0, +\infty)$ for each $i \in \{1, 2\}$ and $c \in I$.*

Proof. Define the function $\bar{\psi}: \mathbb{R} \rightarrow (0, +\infty)$ by

$$\bar{\psi}(x) = \begin{cases} \max\left\{\frac{\sin x^{-1} + 1}{|x|}, |x|\right\} & \text{if } x \neq 0, \\ 1 & \text{if } x = 0. \end{cases}$$

Then, clearly, $\mathcal{C}(\bar{\psi}) = \mathbb{R} \setminus \{0\}$ and $\bar{\psi}[-\delta, 0] = \bar{\psi}[(0, \delta)] = (0, +\infty)$ for each $\delta > 0$. Choose elements $x_1, x_2 \in (a, b)$ and assume that $x_1 < x_2$. Define the function $\psi_1: I \rightarrow (0, +\infty)$ by the formula

$$\psi_1(x) = \begin{cases} \bar{\psi}(x - a) & \text{if } x \in (a, x_1], \\ \bar{\psi}(x - b) & \text{if } x \in [x_2, b), \\ \text{linear,} & \text{on the interval } [x_1, x_2]. \end{cases}$$

Observe that ψ_1 is continuous on I and it is easy to see that $\psi_1[(a, c)] = \psi_1[(c, b)] = (0, +\infty)$ for each $c \in I$. Now, define the function $\psi_2: I \rightarrow (0, +\infty)$ as follows

$$\psi_2 = \frac{f}{\psi_1}.$$

Since f is positive, bounded and continuous on $\text{cl } I$, the function ψ_2 is continuous on I and $\psi_2[(a, c)] = \psi_2[(c, b)] = (0, +\infty)$ for each $c \in I$. Finally, $f = \psi_1 \psi_2$ on I , which completes the proof. $\square$

LEMMA 3.4. *Let $I = (a, b)$ and $y \in (0, 1]$. There is a strong Świątkowski function $\psi: \text{cl } I \rightarrow (0, 1]$ such that $\psi[I] = \psi[I \cap \mathcal{C}(\psi)] = (0, 1]$, $\psi(a) = \psi(b) = y$, $\text{bd } I \subset \mathcal{C}(\psi)$, and $\text{card}(I \setminus \mathcal{C}(\psi)) = 1$.*

P r o o f. Define the function $\bar{\psi}: \mathbb{R} \rightarrow (0, 1]$ by

$$\bar{\psi}(x) = \begin{cases} \min\{1, \sin x^{-1} + |x| + 1\} & \text{if } x \neq 0, \\ 2^{-1} & \text{if } x = 0. \end{cases}$$

Then, clearly, $\bar{\psi} \in \mathcal{S}_s$. Choose elements $x_1, x_2, x_3 \in (a, b)$ and assume that $x_1 < x_2 < x_3$. Define the function $\psi: \text{cl } I \rightarrow (0, 1]$ by the formula

$$\psi(x) = \begin{cases} \bar{\psi}(x - x_2) & \text{if } x \in [x_1, x_3], \\ y & \text{if } x \in \{a, b\}, \\ \text{linear,} & \text{on intervals } [a, x_1] \text{ and } [x_3, b]. \end{cases}$$

One can easily show that the function ψ has all required properties. $\square$

4. Main results

Remark 4.1. Product of two internally cliquish functions is internally cliquish.

P r o o f. If the functions f and g are internally cliquish, then the sets $\text{int } \mathcal{C}(f)$ and $\text{int } \mathcal{C}(g)$ are dense in $\mathbb{R}$. Hence, the set $\text{int } \mathcal{C}(f) \cap \text{int } \mathcal{C}(g)$ is dense in $\mathbb{R}$, too. Moreover, $\text{int } \mathcal{C}(f) \cap \text{int } \mathcal{C}(g) \subset \text{int } \mathcal{C}(fg)$, which proves that the function fg is internally cliquish. $\square$

THEOREM 4.2. *For each function f the following conditions are equivalent:*

- i) *there are internally quasi-continuous functions g_1 and g_2 with $f = g_1 g_2$,*
- ii) *f is a finite product of internally quasi-continuous functions,*
- iii) *f is internally cliquish and the set $[f = 0]$ is simply open.*

P r o o f. The implication i) $\Rightarrow$ ii) is evident.

ii) $\Rightarrow$ iii). Assume that there is $k \in \mathbb{N}$ and there are internally quasi-continuous functions $g_1, \dots, g_k$ such that $f = g_1 \dots g_k$. Since each internally quasi-continuous function is internally cliquish, using Remark 4.1 we obtain that the function f is internally cliquish. Moreover, since each internally quasi-continuous function is quasi-continuous, by Theorem 2.1, $[f = 0]$ is simply open.

iii) $\Rightarrow$ i). Now, assume that the set $[f = 0]$ is simply open and the function f is internally cliquish. Hence, $\text{int } \mathcal{C}(f)$ is dense in $\mathbb{R}$. Let $U = \text{int } \mathcal{C}(f) \setminus \text{bd}[f = 0]$. Observe that the set $\mathbb{R} \setminus U$ is closed. Moreover,

$$\mathbb{R} \setminus U = (\mathbb{R} \setminus \text{int } \mathcal{C}(f)) \cup \text{bd}[f = 0].$$

Since $\mathbb{R} \setminus \text{int } \mathcal{C}(f)$ is boundary and closed, and $[f = 0]$ is simply open, the set $\mathbb{R} \setminus U$ is nowhere dense. Let $\mathcal{J}$ be the family of all components of U . Since $[f = 0]$ is simply open, $J \subset [f = 0]$ or $J \cap [f = 0] = \emptyset$ for each $J \in \mathcal{J}$. So, if there is

PRODUCTS OF INTERNALLY QUASI-CONTINUOUS FUNCTIONS

$J \in \mathcal{J}$ such that $J \cap [f = 0] = \emptyset$, then $f > 0$ on J or $f < 0$ on J . Write the set U as the union of families $\mathcal{I}_1$ and $\mathcal{I}_2$ consisting of the pairwise disjoint compact intervals, such that for each $x \in U$, there are $I_1 \in \mathcal{I}_1$ and $I_2 \in \mathcal{I}_2$ with $x \in \text{int}(I_1 \cup I_2)$.

Fix an interval $J \in \mathcal{J}$ and let $J = (a, b)$. If $J \subset [f = 0]$, then by Lemma 3.1 applied for $\Gamma = +\infty$ and $k = 2$, there are continuous functions $g_{1J}, g_{2J}: J \rightarrow \mathbb{R}$ such that $0 = f|_J = g_{1J}g_{2J}$ and for $i \in \{1, 2\}$

$$g_{iJ}[(a, c)] = g_{iJ}[(c, b)] = \mathbb{R} \quad \text{for each } c \in J. \quad (1)$$

If $J \cap [f = 0] = \emptyset$, then $|f| > 0$ on J . Fix an interval $I \in \mathcal{I}_1 \cup \mathcal{I}_2$ such that $I \subset J$ and let $I = [\alpha, \beta]$. Since $|f|I| > 0$ and f is continuous on I , by Lemma 3.3 there are continuous functions $\psi_{1I}, \psi_{2I}: (\alpha, \beta) \rightarrow (0, +\infty)$ such that $|f| = \psi_{1I}\psi_{2I}$ on (α, β) and for $i \in \{1, 2\}$

$$\psi_{iI}[(\alpha, c)] = \psi_{iI}[(c, \beta)] = (0, +\infty) \quad \text{for each } c \in (\alpha, \beta). \quad (2)$$

Now, define functions $\psi_{1J}, \psi_{2J}: J \rightarrow \mathbb{R}$ as follows:

$$\psi_{1J}(x) = \begin{cases} \psi_{1I}(x) & \text{if } x \in \text{int } I \text{ and } I \in \mathcal{I}_1, \\ -\psi_{1I}(x) & \text{if } x \in \text{int } I \text{ and } I \in \mathcal{I}_2, \\ 1 & \text{otherwise,} \end{cases}$$

$$\psi_{2J}(x) = \begin{cases} \psi_{2I}(x) \cdot \text{sgn } f(x) & \text{if } x \in \text{int } I \text{ and } I \in \mathcal{I}_1, \\ -\psi_{2I}(x) \cdot \text{sgn } f(x) & \text{if } x \in \text{int } I \text{ and } I \in \mathcal{I}_2, \\ f(x) & \text{otherwise.} \end{cases}$$

Then, clearly, $f|_J = \psi_{1J}\psi_{2J}$. By condition (2) and since $|f| > 0$ on J , for $i \in \{1, 2\}$

$$\psi_{iJ}[(a, c)] = \psi_{iJ}[(c, b)] = \mathbb{R} \setminus \{0\} \quad \text{for each } c \in J. \quad (3)$$

ψ_{1J} and ψ_{2J} are internally quasi-continuous on J . Fix $x \in J$. If there is $I \in \mathcal{I}_1 \cup \mathcal{I}_2$ such that $x \in \text{int } I$, then, since ψ_{1I} and ψ_{2I} are continuous on I , the functions ψ_{1J} and ψ_{2J} are internally quasi-continuous at x . In another case, there are $I_1 \in \mathcal{I}_1$ and $I_2 \in \mathcal{I}_2$ such that $x \in \text{bd } I_1 \cap \text{bd } I_2$. Since $\psi_{1J}(x) = 1$ and $\psi_{1J} = \psi_{1I}$ is positive and continuous on $\text{int } I_1$, using (2), we clearly obtain that ψ_{1J} is internally quasi-continuous at x . Moreover, $\psi_{2J}(x) = f(x)$ and ψ_{2J} is continuous on $\text{int } I_1$, it has the same sign as the function f on I_1 . (Recall that f does not change its sign on J .) So, by (2), ψ_{2J} is internally quasi-continuous at x , too.

Further, we define functions $g_1, g_2: \mathbb{R} \rightarrow \mathbb{R}$ as follows:

$$g_1(x) = \begin{cases} g_{1J}(x) & \text{if } x \in J, J \in \mathcal{J} \text{ and } J \subset [f = 0], \\ \psi_{1J}(x) & \text{if } x \in J, J \in \mathcal{J} \text{ and } J \cap [f = 0] = \emptyset, \\ 1 & \text{otherwise,} \end{cases}$$

$$g_2(x) = \begin{cases} g_{2J}(x) & \text{if } x \in J, J \in \mathcal{J} \text{ and } J \subset [f = 0], \\ \psi_{2J}(x) & \text{if } x \in J, J \in \mathcal{J} \text{ and } J \cap [f = 0] = \emptyset, \\ f(x) & \text{otherwise.} \end{cases}$$

Then, clearly, $f = g_1 g_2$. Finally we will show that functions g_1 and g_2 are internally quasi-continuous.

Fix $i \in \{1, 2\}$ and let $x \in \mathbb{R}$. First, assume that there is $J \in \mathcal{J}$ such that $x \in \text{cl } J$. If $x \in \text{int } J$ then, since g_{iJ} is continuous on J and ψ_{iJ} is internally quasi-continuous on J , the function g_i is internally quasi-continuous at x . So, let $x \in \text{bd } J$. Since $g_i|_J$ is internally quasi-continuous, by (1) or (3), we clearly obtain that g_i is internally quasi-continuous at x .

Assume now that $x \in \mathbb{R} \setminus \bigcup_{J \in \mathcal{J}} \text{cl } J$. In this case, $x \in \mathbb{R} \setminus U$. Since $\mathbb{R} \setminus U$ is nowhere dense and conditions (1) and (3) hold, for each $n \in \mathbb{N}$, there is $J_n \in \mathcal{J}$ such that $J_n \subset (x, x + \frac{1}{n})$ and there is

$$x_n \in J_n \cap \text{int } \mathcal{C}(g_i) \quad \text{with} \quad |g_i(x_n) - g_i(x)| < \frac{1}{n}.$$

Hence there is a sequence $(x_n) \subset \text{int } \mathcal{C}(g_i)$ such that $x_n \rightarrow x$. Consequently, the function g_i is internally quasi-continuous at x . This completes the proof. $\square$

PROPOSITION 4.3. *There is a bounded internally quasi-continuous strong Świątkowski function which cannot be written as the finite product of internally strong Świątkowski functions.*

Proof. Let $C \subset [0, 1]$ be the Cantor ternary set, and let $\mathcal{I}_1$ and $\mathcal{I}_2$ be disjoint families of all components of the set $[0, 1] \setminus C$ such that $C \cup \bigcup \mathcal{I}_1 \cup \bigcup \mathcal{I}_2 = [0, 1]$ and $C = (\text{cl } \bigcup \mathcal{I}_1) \cap (\text{cl } \bigcup \mathcal{I}_2)$. Put $\mathcal{I} = \mathcal{I}_1 \cup \mathcal{I}_2$ and define

$$A = C \setminus \bigcup_{I \in \mathcal{I}} \text{bd } I. \quad (4)$$

Since A is a G_δ -set, then $C \setminus A$ is an F_σ -set, whence there is a sequence (F_n) consisting of closed sets such that

$$C \setminus A = \bigcup_{n \in \mathbb{N}} F_n. \quad (5)$$

Define $F'_0 = \emptyset$. For each $n \in \mathbb{N}$, use Lemma 3.2 two times to construct a sequence of sets (F'_n) and a sequence of families of intervals $(\mathcal{J}'_n)$ such that

$$\mathcal{J}'_n = \mathcal{J}'_{1,n} \cup \mathcal{J}'_{2,n}, \quad (6)$$

$$F'_n = F_n \cup \bigcup_{k < n} \left(F'_k \cup \bigcup_{I \in \mathcal{J}'_k} \text{bd } I \right) \quad (7)$$

and for $j \in \{1, 2\}$,

$$\mathcal{J}'_{j,n} \subset \mathcal{I}_j, \quad (8)$$

PRODUCTS OF INTERNALLY QUASI-CONTINUOUS FUNCTIONS

$$\text{for each } I \in \mathcal{I}_j, \text{ if } F'_n \cap \text{bd } I \neq \emptyset, \text{ then } I \in \mathcal{J}'_{j,n}, \quad (9)$$

$$\begin{aligned} &\text{for each } c \in F'_n, \text{ if } c \text{ is a right-hand (left-hand) limit point of } C, \\ &\text{then } c \text{ is a right-hand (left-hand) limit point of the union } \bigcup \mathcal{J}'_{j,n}, \end{aligned} \quad (10)$$

$$\text{cl} \bigcup \mathcal{J}'_{j,n} \subset F'_n \cup \bigcup_{J \in \mathcal{J}'_{j,n}} \text{cl } J. \quad (11)$$

Observe that by (11), for each $k < n$, the set $F'_k \cup \bigcup_{I \in \mathcal{J}'_k} \text{bd } I$ is closed. So, by (7), the set F'_n is also closed and $F'_n \subset C \setminus A$. Fix an interval $I \in \mathcal{I}$. Using Lemma 3.4, we construct a strong Świątkowski function $f_I[I] = f_I[I \cap \mathcal{C}(f_I)] = (0, 1]$, $f_I(\inf I) = f_I(\sup I) = 2^{-1}$, $\text{bd } I \subset \mathcal{C}(f_I)$, and $\text{card}(I \setminus \mathcal{C}(f_I)) = 1$. Put

$$n_I = \min \{n \in \mathbb{N} : I \in \mathcal{J}'_n\},$$

and observe that by (9), $\bigcup_{n \in \mathbb{N}} \bigcup \mathcal{J}'_n = [0, 1] \setminus C$, whence n_I is well defined.

Now, define the function $f: \mathbb{R} \rightarrow [-2^{-1}, 2^{-1}]$ by the formula:

$$f(x) = \begin{cases} (-1)^j 2^{-n_I} f_I(x) & \text{if } x \in \text{cl } I \text{ and } I \in \mathcal{I}_j, j \in \{1, 2\}, \\ 0 & \text{otherwise.} \end{cases}$$

Clearly, f is bounded and $A \subset [f = 0]$. First, we will show that $A \subset \mathcal{C}(f)$.

Take an $x_0 \in A$ and let $\varepsilon > 0$. Choose $n_0 \in \mathbb{N}$ such that $2^{-n_0} < \varepsilon$ and put $\delta = \text{dist}(\text{cl} \bigcup \mathcal{J}'_{n_0}, x_0)$. Since by (11), (7), (4), and (5),

$$\begin{aligned} A \cap \text{cl} \bigcup \mathcal{J}'_{j,n_0} &\subset (A \cap F'_{n_0}) \cup \left(A \cap \bigcup_{J \in \mathcal{J}'_{j,n_0}} \text{cl } J \right) \\ &\subset \left(A \cap \bigcup_{n \leq n_0} F'_n \right) \cup \left((C \setminus \bigcup_{I \in \mathcal{I}} \text{bd } I) \cap \bigcup_{J \in \mathcal{I}} \text{cl } J \right) = \emptyset, \end{aligned}$$

we have $x_0 \notin \text{cl} \bigcup \mathcal{J}'_{j,n_0}$ and $\delta > 0$.

Observe that by (10), $F'_{n_0} \subset \text{cl} \bigcup \mathcal{J}'_{n_0}$. If $|x - x_0| < \delta$, then $x \notin \text{cl} \bigcup \mathcal{J}'_{n_0}$, whence

$$|f(x) - f(x_0)| = |f(x)| \leq 2^{-n_0} < \varepsilon.$$

So, $x_0 \in \mathcal{C}(f)$.

Now, we will prove that

$$\forall_{n \in \mathbb{N}} \forall_{\delta > 0} \left(x \in F'_n \setminus \{\sup I : I \in \mathcal{I}\} \Rightarrow f[(x - \delta, x) \cap \mathcal{C}(f)] \supset [-2^{-n}, 2^{-n}] \right). \quad (12)$$

Let $n \in \mathbb{N}$, $\delta > 0$ and $x \in F'_n \setminus \{\sup I : I \in \mathcal{I}\}$. Then for $j \in \{1, 2\}$, by (10), there is $I_j \in \mathcal{J}'_{j,n}$ with $I_j \subset (x - \delta, x)$. Notice that $\max\{n_{I_1}, n_{I_2}\} \leq n$. So,

$$f[(x - \delta, x) \cap \mathcal{C}(f)] \supset f[I_1 \cap \mathcal{C}(f)] \cup f[I_2 \cap \mathcal{C}(f)] \supset [-2^{-n}, 2^{-n}] \setminus \{0\}.$$

Since $x \notin \{\sup I : I \in \mathcal{I}\}$, we have $(x - \delta, x) \cap A \neq \emptyset$ and finally

$$f[(x - \delta, x) \cap \mathcal{C}(f)] \supset [-2^{-n}, 2^{-n}].$$

Similarly, we can prove that

$$\bigvee_{n \in \mathbb{N}} \bigvee_{\delta > 0} \left(x \in F'_n \setminus \{\inf I : I \in \mathcal{I}\} \Rightarrow f[(x, x + \delta) \cap \mathcal{C}(f)] \supset [-2^{-n}, 2^{-n}] \right).$$

Now, we will show that $f \in \dot{\mathcal{S}}_s$. Let $c, d \in \mathbb{R}$, $c < d$, and $y \in I(f(c), f(d))$. Without loss of generality, we can assume that $c, d \in [0, 1]$ and $f(c) < f(d)$. If $c, d \in \text{cl } I$ for some $I \in \mathcal{I}$, then since $f_I \in \dot{\mathcal{S}}_s$, there is $x_0 \in (c, d) \cap \mathcal{C}(f)$ with $f(x_0) = y$. So, assume that the opposite case holds.

Assume that $y \geq 0$. (The case $y < 0$ is analogous.) Then $f(d) > 0$, whence $d \notin A$. We consider two cases.

Case 1. $d \notin \bigcup_{n \in \mathbb{N}} F'_n$ or $d \in \{\sup I : I \in \mathcal{I}\}$.

Then there is $I \in \mathcal{I}$ such that $d \in \text{cl } I$ and $c \notin \text{cl } I$. If $y \in I(f(\inf I), f(d))$, then, since $f_I \in \dot{\mathcal{S}}_s$, there is $x_0 \in (\inf I, d) \cap \mathcal{C}(f) \subset (c, d) \cap \mathcal{C}(f)$ with $f(x_0) = y$.

Now, let $y \in [0, f(\inf I)]$. By (7), since $I \in \mathcal{J}'_{n_I}$, we have $\inf I \in F'_{n_I+1}$. By (12),

$$y \in [0, f(\inf I)] \subset [-2^{-n_I-1}, 2^{-n_I-1}] \subset f[(c, \inf I) \cap \mathcal{C}(f)].$$

So, there is $x_0 \in (c, \inf I) \cap \mathcal{C}(f) \subset (c, d) \cap \mathcal{C}(f)$ with $f(x_0) = y$.

Case 2. $d \in \bigcup_{n \in \mathbb{N}} F'_n \setminus \{\sup I : I \in \mathcal{I}\}$.

Then, $d \in F'_n \setminus F'_{n-1}$ for some $n \in \mathbb{N}$. By (12),

$$y \in [0, f(d)] \subset [-2^{-n}, 2^{-n}] \subset f[(c, d) \cap \mathcal{C}(f)].$$

Consequently, there is $x_0 \in (c, d) \cap \mathcal{C}(f)$ with $f(x_0) = y$. So, $f \in \dot{\mathcal{S}}_s$.

Observe now that $\mathbb{R} \setminus \mathcal{C}(f) = (C \setminus A) \cup \bigcup_{I \in \mathcal{I}} (I \setminus \mathcal{C}(f_I))$. However, $\text{card}(I \setminus \mathcal{C}(f_I)) = 1$ for each $I \in \mathcal{I}$. So, the set $\mathbb{R} \setminus \mathcal{C}(f)$ is nowhere dense. Hence and since each strong Świątkowski function is quasi-continuous, the function f is internally quasi-continuous.

To complete the proof, suppose that there is $k \in \mathbb{N}$ and there are functions $g_1, \dots, g_k \in \dot{\mathcal{S}}_{si}$ such that $f = g_1 \dots g_k$ on $\mathbb{R}$. Then, $\text{sgn} \circ f = \text{sgn} \circ (g_1 \dots g_k)$. Let $(a, b) \subset [0, 1]$ be an interval in which the function f changes its sign. Then, at least one of the functions $g_1, \dots, g_k$, say g_1 , has the same property. Since $g_1 \in \dot{\mathcal{S}}_{si}$, there is $x_0 \in (a, b) \cap \text{int } \mathcal{C}(g_1)$ such that $g_1(x_0) = 0$. Note that $[f = 0] \cap [0, 1] = A$, whence $g(x_1) \neq 0$ for some $x_1 \in (a, b) \cap \text{int } \mathcal{C}(g_1) \cap (C \setminus A)$. Since g_1 is continuous at x_1 , there is an open interval (a_1, b_1) such that $g_1[(a_1, b_1)] \cap [g_1 = 0] = \emptyset$ and f changes its sign in (a_1, b_1) . Hence at least one of the functions $g_2, \dots, g_k$, say g_2 , changes its sign in (a_1, b_1) , too. Observe that $g_2 \in \dot{\mathcal{S}}_{si}$. Proceeding as above, after k steps, we obtain that there is an open interval J in which the function f changes its sign and $g_i[J] \cap [g_i = 0] = \emptyset$ for each $i \in \{1, \dots, k\}$, a contradiction.

Consequently, the function f cannot be written as the finite product of internally strong Świątkowski functions. $\square$

PRODUCTS OF INTERNALLY QUASI-CONTINUOUS FUNCTIONS

Finally, we would like to present the problem.

PROBLEM 4.4. Characterize the products of internally strong Świątkowski functions.

REFERENCES

- [1] BISWAS, N.: *On some mappings in topological spaces*, Bull. Calcutta Math. Soc. **61** (1969), 127–135.
- [2] CEDER, J. G.: *On factoring a function into a product of Darboux functions*, Rend. Circ. Mat. Palermo (2) **31** (1982), 16–22.
- [3] CEDER, J. G.: *A necessary and sufficient condition for a Baire α function to be a product of two Darboux Baire α functions*, Rend. Circ. Mat. Palermo (2) **36** (1987), 78–84.
- [4] GRANDE, Z.: *Sur le fonctions cliquish*, Časopis Pěst. Mat. **110** (1985), 225–236.
- [5] KEMPISTY, S.: *Sur les fonctions quasicontinues*, Fund. Math. **19** (1932), 184–197.
- [6] MALISZEWSKI, A.: *On the limits of strong Świątkowski functions*, Zeszyty Nauk. Politech. Łódz. Mat. **27** (1995), 87–93.
- [7] MALISZEWSKI, A.: *Darboux Property and Quasi-Continuity. A Uniform Approach*. WSP, Szupsk, 1996.
- [8] MARCINIAK, M.—SZCZUKA, P.: *On internally strong Świątkowski functions* (preprint).
- [9] MARCUS, S.: *Sur la représentation d'une fonction arbitraire par des fonctions jouissant de la propriété de Darboux*, Trans. Amer. Math. Soc. **95** (1960), 484–494.
- [10] SZCZUKA, P.: *Products of strong Świątkowski functions*, J. Appl. Anal. **12** (2006), 129–145.
- [11] THIELMAN, H. P.: *Types of functions*, Amer. Math. Monthly **60** (1953), 156–161.

Received October 12, 2012

*Institute of Mathematics
Kazimierz Wielki University
pl. Weyssenhoffa 11
PL-85-072 Bydgoszcz
POLAND
E-mail: paulinaszczuka@wp.pl
marmac@ukw.edu.pl*